

Hani H. Abdel-Nabi, M.D., Ph.D., F.A.C.N.P.

Director of Nuclear Medicine Residency Program
State University of New York at Buffalo
Department of Nuclear Medicine
105 Parker Hall
3435 Main Street
Buffalo, NY 14214-3007

Dr. Hani Abdel-Nabi is a Professor of Nuclear Medicine in the Department of Nuclear Medicine, State University of New York at Buffalo. After earning his medical degree from the University of Alexandria School of Medicine (Egypt), Dr. Nabi completed a residency in Radiation Oncology at the same institution.

In 1979, Dr. Nabi came to the Ohio State University on a Fullbright Scholarship where he earned a Ph.D. in Radiation Biology. He also joined the Department of Radiology at OSU as a resident in Nuclear Medicine, then a fellow from 1979-1983.

Dr. Nabi is a diplomate of the American Board of Nuclear Medicine. His research interests are mostly in cancer detection and treatment, specifically with radiolabeled monoclonal antibodies and more recently positron emission tomography. Dr. Nabi is author of 59 journal articles, 5 book chapters, and 135 abstracts.

He is currently the director of the Residency Training Program at SUNY and is active in academic medicine as well as in professional organizations, notably the Society of Nuclear Medicine.

Immunoscintigraphy of Indeterminate Breast Lesions

Breast Cancer is the most frequently diagnosed neoplasia in American women and the second leading cause of cancer mortality. The incidence of breast cancer increases sharply for American women in their early 40's. It then levels off after 45 years and increases again after age 55. For women 70 years old, the yearly incidence is three times that of those age 50.

Mortality from breast cancer is strongly influenced by the stage of disease at detection. An overall 8.1% decrease in breast cancer mortality was reported for 1989-1992 by the Surveillance, Epidemiology and End Results Program. Part of this decrease can be attributed to screening mammography which is the most effective approach to the early detection of breast cancer.

As a result of the widespread use of screening mammography (national rate is currently 71.2%), more abnormalities such as mass and or microcalcifications are discovered yearly. Although many of these abnormalities lack the characteristic features of malignancies, up to 700,000 women undergo biopsies for suspicious mammograms yearly, and only one out of seven biopsies prove to be cancer. Over the past decades, efforts have been directed towards the development of alternative methods that would lead to better characterization of malignant lesions.

Ultrasonography has been proven to be of benefit in separating benign simple cysts, which are unlikely to be malignant, from those complex masses which may require tissue diagnosis. The role of both magnetic resonance and positron emission tomography with ^{18}F -Fluorodeoxyglucose is still under investigation. The lipophilic agent, sestamibi has been advocated in patients with indeterminate mammograms and dense dysplastic breasts. Results of the multinational trial published recently demonstrated an accuracy of approximately 85% for breast masses larger than 1.0 cm in diameter. Recent evidence points to the presence of false-positive findings due primarily to an interaction between sestamibi and benign hyperplasia, or fibroadenomas. False-positive rates of up to 35% have been reported.

The recent development of monoclonal antibodies with high reactivity with tumor-specific or associated antigens offers a new avenue towards improved tumor targeting.

Arcitumomab, a technetium- $^{99\text{m}}$ based monoclonal antibody Fab' fragments have been evaluated in patients with breast lesions as part of a multinational center trial. For a total of 139 patients, immunoscintigraphy had an overall sensitivity of 82% and a specificity of 90%. When patients with indeterminate suspicious mammograms (BiRads 3 and 4) were evaluated, Arcitumomab had a sensitivity of 62% and a specificity of 97%, which was significantly higher than mammography (97% vs. 75%, $p < 0.01$)

Using these results to compute likelihood ratios for having or not having cancer, a positive Arcitumomab increased a patient's odds of having cancer 7-fold, while a negative imaging result decreased these odds 5-fold.

As expected, tumor detection rates were influenced by the size of the lesion, varying 60% for lesions below 1.0 cm, to 92% for lesions greater than 1.1 cm. This limitation can potentially be minimized or negated by the use of positron emitting radioisotopes labeled to the monoclonal antibody fragments, Copper-64, Iodine-124, Technetium-94m or Fluorine-18 are potential candidates that require further exploration. Features specific to each of these radioisotopes will be discussed.

References:

1. Cardenosa G. (1977). Breast Cancer and mammography: an overview. In *Breast Imaging Companion*. New York: Lippincott-Raven
2. Newman L. (1977). National Institutes of Health Consensus development conferences statements, breast cancer screening for women ages 40-49. In *1998 Medical Outcomes & Guidelines Sourcebook*. New York: Faulkner & Gray.
3. Pederson L. & Trigg J. (1995). Finding and reacting to a lump. *Breast Cancer, A Family Survival Guide*. Westport, Connecticut: Bergin & Garvey.
4. U.S. Department of Health and Human Services, Public Health Service. *Healthy People 2000*, Washington, D.C.: U.S. Government Printing Office.
5. Miller RS, Adelman RW, Espinosa MH, Dorman SA, Smith DH. The early detection of nonpalpable breast carcinoma with needle localization. Experience with 500 patients in a community hospital. *Am Surg* 1992; 58: 193-198.
6. Opie H, Estes NC, Jewell WR, Chang CH, Thomas JA, Estes MA. Breast biopsy for nonpalpable lesions: a worthwhile endeavor? *Am Surg* 1993; 59:490-493.
7. Knutzen AM, Gisvold JJ. Likelihood of malignant disease for various categories of mammographically detected, nonpalpable breast lesions. *Mayo Clin Proc* 1993; 68:454-460.
8. Rosenberg RD, Lando JF, Hunt WC et al. The New Mexico Mammography Project. Screening mammography performance in Albuquerque, New Mexico, 1991 to 1993. *Cancer* 1996; 78:1731-1739.
9. Khalkhali I, Cutrone JA, Mena I. Scintimammography: the complementary role of Tc-99m sestamibi prone breast imaging for the diagnosis of breast carcinoma. *Radiology* 1995;196:421-426.
10. Scopinaro F, Schillaci O, Ussof W, et al. A three center study on the diagnostic accuracy of 99mTc-MIBI scintimammography. *Anticancer Res* 1997;17:1631-1634.
11. Palmedo H, Biersack HJ, Lastoria S, Maublant J, Prats E, Stegner HE, Bourgeois P, Hustinx R, Hilson AJW, Bischof-Delaloye A. Scintimammography with technetium-99m methoxyisobutylisonitrile: results of a prospective European multicentre trial. *Eur J Nucl Med* 1998; 25:375-385.
12. Tiling R, Khalkhali I, Sommer H, et al. Limited value of scintimammography and contrast-enhanced MRI in the evaluation of microcalcification detected by mammography. *Nucl Med Commun* 1998; 19:55-62.
13. Goldenberg DM, DeLand F, Kim E, et al. Use of radiolabeled antibodies to carcinoembryonic antigen for the detection and localization of diverse cancers by external photoscanning. *N Engl J Med* 1978; 298:1384-1388.
14. Bocker W, Schweikhart G, Pollow K, et al. Immunohistochemical demonstration of carcinoembryonic antigen (CEA) in 120 mammary carcinomas and its correlation with tumor type, grading, staging plasma-CEA, and biochemical receptor status. *Pathol Res Pract* 1985; 180:490-497.
15. Nabi HA. Antibody imaging in breast cancer. *Semin Nucl Med* 1997;27:30-39.
16. Tolmos J, Cutrone JA, Wang B, et al. Scintimammographic analysis of nonpalpable breast lesions previously identified by conventional mammography. *J Natl Cancer Inst* 1998; 90:846-849.
17. Goldenberg DM, Nabi HA. Breast Cancer Imaging with Radiolabeled Antibodies. *Semin Nucl Med* 1999; 29(1):41-48.
18. Goldenberg DM, Abdel-Nabi H, Serafini A, Seldin D, Barron B, Line B, Wegner WA. The role of carcinoembryonic antigen immunoscintigraphy in the diagnosis of breast carcinoma. *Breast Cancer Research & Treatment*, 21st annual San Antonio Breast Cancer Symposium 1998; 50(3):316 #501.